

Spin-resolved Photoemission Studies of Gd(0001)

A.V. Fedorov #

T. Valla

P.D. Johnson

M. Weinert

/Brookhaven/

P.B. Allen /Stony Brook/

now at ALS

Spin Resolved Photoemission on Gd

- ★ Surface Curie temperature ($T_{\text{surf}} \sim T_{\text{bulk}} + 60\text{K}$)
- ★ Canted magnetic moments
- ★ Stoner splitting vs. spin-mixing
- ★ Electron-magnon coupling 

Plan


- ★ Photoemission, self energies, scattering rates,etc.
- ★ Micro-Mott spin polarimeter, achieving the high resolution
- ★ Scattering channels for the spin majority and minority photo-holes
- ★ Surface Curie temperature

Angle Resolved Photoemission /band structure mapping/


Experiment

Excitation Radiation

- photon energy
- polarization
- angle of incidence




- ### Photoelectrons
- kinetic energy
 - emission angle
 - polarization



Data



Energy Distribution Curves (photocurrent vs. kinetic energy) measured at certain emission angle



Angle Resolved Photoemission /spectral function and self energy/



$$A(\kappa, \omega) = \frac{1}{\pi} \frac{\text{Im}\Sigma(\kappa, \omega)}{[\omega - \varepsilon_\kappa - \text{Re}(\kappa, \omega)]^2 + [\text{Im}\Sigma(\kappa, \omega)]^2}$$

$\text{Im}\Sigma(\kappa, \omega) \sim \frac{1}{T}$ and gives the scattering rate /e-ph, e-e, e-imp, **e-magnon**/



High-efficiency retarding-potential Mott polarization analyzer

$$V_{so} = \frac{1}{2m^2c^2} \frac{1}{r} \frac{dV}{dr} \propto \rho$$




G. C. Burnett, T. J. Monroe, and F. B. Dunning

Department of Physics and the Rice Quantum Institute, Rice University, P.O. Box 1892, Houston, Texas 77251

(Received 17 January 1994; accepted for publication 14 March 1994)

/Rev. Sci. Instrum. **65**, 1893/



$$P = \frac{1}{S} \frac{\sqrt{I_L^+ I_R^-} - \sqrt{I_L^- I_R^+}}{\sqrt{I_L^+ I_R^-} + \sqrt{I_L^- I_R^+}}$$

$$A = \frac{\sqrt{I_L^+ I_L^-}}{\sqrt{I_R^+ I_R^-}} \quad \text{Instrumental asymmetry}$$


$$I^\uparrow = \langle I \rangle (1+P)$$

$$I^\downarrow = \langle I \rangle (1-P)$$


$$\langle I \rangle = \frac{I_L^+ + I_L^- + I_R^+ + I_R^-}{4}$$

$$\eta = S_{\text{eff}}^2 / I_0$$

Spin polarimeter in BNL




Energy resolution: {Aperture/ \varnothing MCP} \times {PE/10}






Aperture plate




Channeltrons




90° deflector

Photoelectron Spectrometer

/200-mm hemispherical analyzer from Scienta/




Test results:
"in-plane" polarization in
ferromagnetic iron films




Resolution: 50 meV / PE=10eV/
Sherman function: 0.15


Surface state in Gd(0001)



D. Li et al., PRB **51**, 13895



E. Weschke et al., PRL **77**, 3415



Sample Preparation:


Epitaxial films of Gd on Mo(110)
Deposition at 200÷300 K
Annealing at ~800 K
Result: Gd(0001)

Sample lifetime ~ 48 Hours


Temperature range 20 K÷3000 K

Base pressure 2×10^{-11} mbar

Angle-resolved spin-integrated data
/suggests Stoner-like behavior/



Spin-resolved data



Temperature-dependent shift
of spin-majority peak agrees
with Stoner behavior


Spin-majority peak shows considerable
broadening with temperature

Surface state is not 100%
polarized even at 20K

Relative intensity of spin-minority
component increases with temperature

Spectra were measured at six different temperatures:
20K, 80K, 98K, 117K, 140K, 160K

Width of spin-majority peak vs. temperature



Broadening of majority peak reflects mostly phonon contribution to the line-width

"Spin-resolved" electron-phonon coupling constants:
 $\lambda \uparrow \sim 0.73$ $\lambda \downarrow \sim 0.31$ /P. Allen/

"Spin-averaged" constant: 0.4
 /H.L. Skriver and I. Mertig, PRB **41**, 6553



Experiment: $\lambda \sim 0.9$

$$\text{Im} \sum_{e-ph}(T, \omega) = \pi \int d\nu \alpha^2 F(\nu) [1 + 2n(\nu) + f(\nu + \omega) - f(\omega - \nu)]$$

Debye model: $\alpha^2 F(w) = \lambda (\omega / \Omega_m)^2$ $1/\tau \approx \text{const} + \pi \lambda k_B T$


Extra broadening of spin-minority peak was detected at all temperatures

Least-square fits to Lorentzian + Fermi edge



Intrinsic width of the peak should not depend on the orientation of magnetic moment in domains

Broader minority spin channel indicates electron-magnon mechanism



P.B. Allen, PRB **63**, 214410 (2001)

$$\frac{1}{\tau^{\downarrow}} \approx \frac{P(\uparrow)m^*}{S} \left(\frac{2JSa}{\eta} \right)^2$$




$$\frac{1}{\tau^{\downarrow}} \approx 0.095 \text{ eV}$$

$$\frac{1}{\tau^{\uparrow}} \approx 0.014 \text{ eV}$$


At low T majority spin hole decays primarily via lattice excitations, minority spin hole primarily via spin excitations

	Majority	Minority
Phonons	46 meV	10 meV
Magnons	14 meV	95 meV




Surface transition temperature: enhanced or regular?

C.S. Arnold and D.P. Pappas,
PRL 85, 5202 (2000):
Common Curie temperature
for surface and bulk




Present work at ~ 300 K:
Hint for the enhanced T_c



Polarization of the surface state vs. temperature

$$\text{Polarization} = \{n\uparrow - n\downarrow\} / \{n\uparrow + n\downarrow\}$$



Measuring “out-of-plane” component in Gd

